



TITLE:

A note on asymptotic solutions of Hamilton-Jacobi equations(Viscosity Solution Theory of Differential Equations and its Developments)

AUTHOR(S):

FUJITA, Yasuhiro

CITATION:

FUJITA, Yasuhiro. A note on asymptotic solutions of Hamilton-Jacobi equations(Viscosity Solution Theory of Differential Equations and its Developments). 数理解析研究所講究録 2006, 1481: 88-92

ISSUE DATE:

2006-04

URL:

<http://hdl.handle.net/2433/58054>

RIGHT:

A note on asymptotic solutions of Hamilton-Jacobi equations

Yasuhiro FUJITA (藤田 安啓)

University of Toyama, 930-8555 Toyama, Japan

(富山大学 理学部)

yfujita@sci.toyama-u.ac.jp

This is a survey of my result [10]. In this talk, we consider the viscosity solutions of the Cauchy problem

$$(1) \quad u_t + \alpha x \cdot Du + H(Du) = f \quad \text{for } (x, t) \in \mathbb{R}^N \times (0, \infty),$$

$$(2) \quad u|_{t=0} = \phi \quad \text{for } x \in \mathbb{R}^N,$$

where α is a positive constant. Our goal is to investigate convergence rates of $u(t, x)$ to the stationary state as $t \rightarrow \infty$. We assume the following:

$$(A1) \quad H, f, \phi \in C(\mathbb{R}^N).$$

$$(A2) \quad H \text{ is convex on } \mathbb{R}^N.$$

$$(A3) \quad \lim_{|x| \rightarrow \infty} \frac{H(x)}{|x|} = \infty.$$

We denote by L the convex conjugate of H defined by

$$L(x) = \sup \{ z \cdot x - H(z) \mid z \in \mathbb{R}^N \}.$$

Then, L satisfies (A2) and (A3) in place of H . Furthermore, we assume that there is a convex function ℓ on \mathbb{R}^N satisfying

$$(A4) \quad \lim_{|x| \rightarrow \infty} (L(x) - \ell(x)) = \infty.$$

$$(A5) \quad \inf \{ f(x) + \ell(-\alpha x) \mid x \in \mathbb{R}^N \} > -\infty.$$

$$(A6) \quad \inf \left\{ \phi(x) + \frac{1}{\alpha} \ell(-\alpha x) \mid x \in \mathbb{R}^N \right\} > -\infty.$$

Now, we introduce several notations.

$$c := \min \left\{ f(x) + L(-\alpha x) \mid x \in \mathbb{R}^N \right\}, \quad f_c(x) := f(x) - c,$$

$$Z := \{x \in \mathbb{R}^N \mid f_c(x) + L(-\alpha x) = 0\},$$

$$\mathcal{C}(x, T) = \{X \in AC([0, T]) \mid X(0) = x\},$$

$$\mathcal{C}(x, y, T) = \{X \in \mathcal{C}(x, T) \mid X(T) = y\},$$

$$d(x, y) = \inf \left\{ \int_0^T [f_c(X(t)) + L(-\alpha X(t) - \dot{X}(t))] dt \mid T > 0, X \in \mathcal{C}(x, y, T) \right\},$$

$$\psi(x) = \inf \left\{ \int_0^T [f_c(X(t)) + L(-\alpha X(t) - \dot{X}(t))] dt + \phi(X(T)) \mid T > 0, X \in \mathcal{C}(x, T) \right\},$$

$$v(x) = \min_{z \in Z} (d(x, z) + \psi(z)).$$

The following propositions were proved in [11] (see also the paper of Professor Hitoshi Ishii in this volume).

Proposition 1. There is the unique viscosity solution $u \in C(\mathbb{R}^N \times [0, \infty))$ of (1)-(2) satisfying for any $T > 0$

$$(3) \quad \lim_{r \rightarrow \infty} \inf \left\{ u(x, t) + \frac{1}{\alpha} L(-\alpha x) \mid (x, t) \in (\mathbb{R}^N \setminus B(0, r)) \times [0, T] \right\} = \infty,$$

where $B(a, r) = \{x \in \mathbb{R}^N \mid |x - a| \leq r\}$ for $a \in \mathbb{R}^N$ and $r > 0$. \square

Proposition 2. For the unique viscosity solution $u \in C(\mathbb{R}^N \times [0, \infty))$ of (1)-(2) satisfying (3), we have

$$(4) \quad \lim_{t \rightarrow \infty} \max_{x \in B(0, R)} |u(x, t) - (ct + v(x))| = 0 \quad \text{for } R > 0. \quad \square$$

Note that by the stability theorem of viscosity solutions, v is a viscosity solution of the equation

$$(5) \quad c + \alpha x \cdot Dv + H(Dv) = f \quad \text{for } x \in \mathbb{R}^N.$$

Next, we consider the convergence rate of (4). First, we consider the case such that the convergence rate of (4) is faster than $e^{-\theta t}$ for some constant $\theta > 0$. Besides (A1) \sim (A6), we assume the following:

(A7) $H \geq 0$ in \mathbb{R}^N with $H(0) = 0$.

(A8) $f \geq 0$ in \mathbb{R}^N with $f(0) = 0$, and there exists a constant $\theta > 0$ such that

$$\theta \int_0^\infty f(xe^{-\alpha t}) dt \leq f(x) \quad \text{for } x \in \mathbb{R}^N.$$

(A9) There exists a constant $m > 0$ such that

$$0 \leq \phi(x) \leq m w(x) \quad \text{for } x \in \mathbb{R}^N,$$

where $w \in C(\mathbb{R}^N)$ is a subsolution of

$$\alpha x \cdot Dv(x) + H(Dv(x)) = f(x) \quad \text{in } \mathbb{R}^N,$$

and satisfies the following inequality for a constant $\lambda > 0$:

$$0 \leq \lambda w(x) \leq f(x) \quad \text{for } x \in \mathbb{R}^N.$$

Example 1. Let f be a nonnegative and convex function on \mathbb{R}^N with $f(0) = 0$. Then, f satisfies (A8) for $\theta = \alpha$.

Example 2. Let G be a nonnegative and convex function on \mathbb{R}^N with $G(0) = 0$. Assume that there exist constants δ_1, δ_2 ($0 < \delta_1 < \delta_2$) and $f \in C(\mathbb{R}^N)$ such that

$$\delta_1 G(x) \leq f(x) \leq \delta_2 G(x) \quad \text{for } x \in \mathbb{R}^N.$$

Then, f satisfies (A8) for $\theta = \alpha \delta_1 / \delta_2$.

Example 3. Assume that there are constants $p \in (1, \infty)$ and $a \in (0, 1)$ such that

$$a \left(\alpha |x|^p + H(|x|^{p-2} x) \right) \leq f(x) \quad \text{for } x \in \mathbb{R}^N.$$

Then, as a function $\phi \in C(\mathbb{R}^N)$ of (A9), we can take any one satisfying

$$0 \leq \phi(x) \leq k |x|^p \quad \text{for } x \in \mathbb{R}^N,$$

where $k > 0$ is a constant.

Theorem 3. Assume (A1)-(A9). Let $u \in C(\mathbb{R}^N \times [0, \infty))$ be the unique viscosity solution of (1)-(2) satisfying (3). Then, we have $c = 0$, $Z \ni 0$, and,

$$(6) \quad -v(x)e^{-\theta t} \leq u(x, t) - v(x) \leq mv(x)e^{-\theta t} \quad \text{for } (x, t) \in \mathbb{R}^N \times [0, \infty).$$

Finally, we give an example, which shows that there is the case such that the convergence rate of (4) is not faster than t^{-1} .

Example 4. Let $H(x) = |x|^p/p$ for some constant $p > 1$. Then, $L(x) = |x|^q/q$, where $(1/p) + (1/q) = 1$. For $r > 0$, let

$$f(x) = -\frac{\alpha^q}{q} \min\{|x|^q, r^q\} \quad \text{for } x \in \mathbb{R}^N.$$

Let $\phi \in C(\mathbb{R}^N)$ be a function satisfying $\phi(x) \geq 0$ for $x \in \mathbb{R}^N$. Then, we have $c = 0$, $Z = B(0, r)$, and,

$$(7) \quad \frac{1}{\alpha} L(-\alpha x)(t+1)^{-1} \leq u(x, t) - v(x) \quad \text{for } (x, t) \in Z \times [0, \infty).$$

References

- [1] O. ALVAREZ, *Bounded-from-below viscosity solutions of Hamilton-Jacobi equations*, Differential Integral Equations 10 (1997), no. 3, 419–436.
- [2] G. BARLES, *Solutions de viscosité des équations de Hamilton-Jacobi*, Mathématiques & Applications (Berlin), Vol. 17, Springer-Verlag, Paris, 1994.
- [3] H. BRÉZIS, *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*, North-Holland Mathematics Studies, No. 5. Notas de Matemática (50). North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, 1973.
- [4] M. BARDI AND I. CAPUZZO-DOLCETTA, *Optimal control and viscosity solutions of Hamilton-Jacobi-Bellman equations*. With appendices by Maurizio Falcone and Pierpaolo Soravia, Systems & Control: Foundations & Applications. Birkhäuser Boston, Inc., Boston, MA, 1997.
- [5] G. BARLES AND P. E. SOUGANIDIS, *On the large time behavior of solutions of Hamilton-Jacobi equations*, SIAM J. Math. Anal. 31 (2000), no. 4, 925–939.

- [6] M. G. CRANDALL, H. ISHII, AND P.-L. LIONS, *User's guide to viscosity solutions of second order partial differential equations*, Bull. Amer. Math. Soc. 27 (1992), 1–67.
- [7] A. DAVINI AND A. SICONOLFI, *A generalized dynamical approach to the large time behavior of solutions of Hamilton-Jacobi equations*, preprint, 2005.
- [8] A. FATHI, *Théorème KAM faible et théorie de Mather pour les systèmes lagrangiens*, C. R. Acad. Sci. Paris Sér. I 324 (1997) 1043–1046
- [9] A. FATHI, *Sur la convergence du semi-groupe de Lax-Oleinik*, C. R. Acad. Sci. Paris Sér. I Math. 327 (1998), no. 3, 267–270.
- [10] Y. FUJITA, *Convergence rates of asymptotic solutions to Hamilton-Jacobi equations in Euclidean n space*, preprint.
- [11] Y. FUJITA, H. ISHII, AND P. LORETI, *Asymptotic solutions to Hamilton-Jacobi equations in Euclidean n space*, preprint.
- [12] H. ISHII, *Comparison results for Hamilton-Jacobi equations without growth condition on solutions from above*, Appl. Anal. 67 (1997), no. 3-4, 357–372.
- [13] J.-M. ROQUEJOFFRE, *Convergence to steady states or periodic solutions in a class of Hamilton-Jacobi equations*, J. Math. Pures Appl. (9) 80 (2001), no. 1, 85–104.